

Some results on counting roots of polynomials and the Sylvester resultant.

Michael Monagan and Baris Tuncer

Department of Mathematics, Simon Fraser University, Burnaby, B.C., V5A 1S6, CANADA.

Abstract. We present two results, the first on the distribution of the roots of a polynomial over the ring of integers modulo n and the second on the distribution of the roots of the Sylvester resultant of two multivariate polynomials. The second result has application to polynomial GCD computation and solving polynomial diophantine equations.

Keywords: roots of polynomials, finite fields, the Sylvester resultant, unlucky evaluation points

1 Introduction

Let \mathbb{F}_q denote the finite field with q elements and let \mathbb{Z}_n denote the ring of integers modulo n . Let $E[X]$ denote the expected value of a random variable X and let $\text{Var}[X]$ denote the variance of X .

Let f be a polynomial in $\mathbb{F}_q[x]$ of a given degree $d > 0$ and let X be the number of distinct roots of f . Schmidt proves in Ch. 4 of [9] that $E[X] = 1$ and for $d > 1$, $\text{Var}[X] = 1 - 1/q$. This result has been generalized by Knopfmacher and Knopfmacher in [5] who count distinct irreducible factors of a given degree of f . The two main results presented in this paper are Theorems 1 and 2 below.

Theorem 1 Let $\phi(n) = |\{1 \leq i \leq n : \gcd(i, n) = 1\}|$ denote Euler's totient function. Let X be a random variable which counts the number of distinct roots of a monic polynomial in $\mathbb{Z}_n[x]$ of degree $m > 0$. Then

(a) $E[X] = 1$ and

(b) if $m = 1$ then $\text{Var}[X] = 0$, otherwise $\text{Var}[X] = \sum_{d|n, d \neq n} \frac{d}{n} \phi\left(\frac{n}{d}\right) = \sum_{d|n} \frac{d-1}{n} \phi\left(\frac{n}{d}\right)$.

In particular, if $n = p^k$ where p is a prime number and $k \geq 1$, $\text{Var}[X] = k(1 - 1/p)$.

Theorem 2 Let f, g be polynomials in $\mathbb{F}_q[x, y]$ of the form $f = c_n x^n + \sum_{i=0}^{n-1} \sum_{j=0}^{n-i} c_{ij} x^i y^j$ and $g = d_m x^m + \sum_{i=0}^{m-1} \sum_{j=0}^{m-i} d_{ij} x^i y^j$ with $c_n \neq 0$ and $d_m \neq 0$, thus of total degree n and m respectively. Let X be a random variable that counts the number of $\gamma \in \mathbb{F}_q$ such that $\gcd(f(x, \gamma), g(x, \gamma)) \neq 1$. If $n > 0$ and $m > 0$ then

(a) $E[X] = 1$ and

(b) $\text{Var}[X] = 1 - 1/q$.

Theorems 1 and 2 were found by computation. We give some details on our computations later in the paper. To prove the results we use a generalization of the Inclusion Exclusion principle (Proposition 1) which allows us to determine $E[X]$ and $\text{Var}[X]$ without having explicit formulas for $\text{Prob}[X = k]$.

Before proving these results we connect Theorem 2 with the Sylvester resultant and with polynomial GCD computation and with solving polynomial diophantine equations.

Let F be a field and let A and B be polynomials in $F[x_0, x_1, \dots, x_n]$ with positive degree in x_0 . The Sylvester resultant of A and B in x_0 , denoted $\text{res}_{x_0}(A, B)$, is the determinant of Sylvester's matrix. We gather the following facts about it into Lemma 1 below. Proofs may be found in Ch. 3 of [3]. Note, in the Lemma $\deg A$ denotes the total degree of A .

Lemma 1 Let $R = \text{res}_{x_0}(A, B)$

- (i) R is a polynomial in $F[x_1, \dots, x_n]$ (x_0 is eliminated),
- (ii) $\deg R \leq \deg A \deg B$ (Bezout bound).

For A and B monic in x_0 and $\alpha \in F^n$

- (iii) $\gcd(A(x_0, \alpha), B(x_0, \alpha)) \neq 1 \iff \text{res}_{x_0}(A(x_0, \alpha), B(x_0, \alpha)) = 0$ and
- (iv) $\text{res}_{x_0}(A(x_0, \alpha), B(x_0, \alpha)) = R(\alpha)$.

Properties (iii) and (iv) connect the roots of the resultant with Theorem 2 and 3.

1.1 Polynomial GCD computation and polynomial diophantine equations.

Our motivation comes from the following problems in computer algebra. Let A, B be polynomials in $\mathbb{Z}[x_0, x_1, \dots, x_n]$ and $G = \gcd(A, B)$. Thus $A = G\hat{A}$ and $B = G\hat{B}$ for some polynomials \hat{A} and \hat{B} called the cofactors of A and B . Modular GCD algorithms compute G modulo a sequence of primes p_1, p_2, p_3, \dots and recover the integer coefficients of G using Chinese remaindering. The fastest algorithms for computing G modulo a prime p interpolate G from univariate images. Maple, Magma and Mathematica all currently use Zippel's algorithm (see [11, 4]). Let us write

$$A = \sum_{i=0}^k a_i x_0^i, \quad B = \sum_{i=0}^l b_i x_0^i, \quad \text{and} \quad G = \sum_{i=0}^m c_i x_0^i$$

where the coefficients $a_i, b_i, c_i \in \mathbb{F}_p[x_1, \dots, x_n]$. Zippel's algorithm picks points $\alpha_i \in \mathbb{F}_p^n$, computes monic univariate images of G

$$g_i = \gcd(A(x_0, \alpha_i), B(x_0, \alpha_i)),$$

scales them (details omitted), then interpolates the coefficients $c_i(x_1, \dots, x_n)$ of G from the coefficients of these (scaled) images.

What if $\gcd(\hat{A}(x_0, \alpha_j), \hat{B}(x_0, \alpha_j)) \neq 1$ for some j ? For example, if $\hat{A} = x_0^2 + x_2$ and $\hat{B} = x_0^2 + x_2 + (x_1 - 1)$ then $\gcd(\hat{A}, \hat{B}) = 1$ but $\gcd(\hat{A}(x_0, 1, \beta), \hat{B}(x_0, 1, \beta)) \neq 1$ for all $\beta \in \mathbb{F}_p$. The evaluation points $(1, \beta)$ are said to be *unlucky*. We cannot use the images $\gcd(A(x_0, 1, \beta), B(x_0, 1, \beta))$ to interpolate G . The same issue of unlucky evaluation points arises in our current work in [6] where, given polynomials $a, b, c \in \mathbb{Z}[x_0, x_1, \dots, x_n]$ with $\gcd(a, b) = 1$ we want to solve the diophantine equation $\sigma a + \tau b = c$ for σ and τ in $\mathbb{Z}[x_0, x_1, \dots, x_n]$ by interpolating σ and τ modulo a prime p from univariate images.

What is the maximum number of unlucky evaluation points that can occur? And what is the expected number of unlucky evaluation points? We answer the first question for A and B monic in x_0 . Lemma 1 implies α_j is unlucky if and only if $R(\alpha_j) = 0$ where $R = \text{res}_{x_0}(\hat{A}, \hat{B}) \in \mathbb{F}_p[x_1, \dots, x_n]$. If α_j is chosen at random from \mathbb{F}_p^n then applying the Schwarz-Zippel lemma (see [10]) we have

$$\text{Prob}[R(\alpha_j) = 0] \leq \frac{\deg R}{p}.$$

Applying Lemma 1(ii) we have $\deg R \leq \deg \widehat{A} \deg \widehat{B} \leq \deg A \deg B$. So if the algorithm needs, say, t images to interpolate G modulo p , then we can avoid unlucky evaluation points with high probability if we pick $p \gg t \deg A \deg B$.

But this is an upper bound – a worst case bound for the GCD algorithm. Researchers in computer algebra have observed that unlucky evaluation points are rare in practice and that we “never see them” when testing algorithms on random inputs. Theorems 2 and 3 give first results on the distribution of unlucky evaluation points. In particular, for coprime \widehat{A} and \widehat{B} of positive degree, Theorem 3 (page 11) implies $\text{Prob}[\alpha_j \text{ is unlucky}] < 1/p$.

2 Results and Proofs

Given a set U and the finite collection of sets $\Gamma = \{A_i, i = 0, \dots, n-1\}$ where each $A_i \subseteq U$, let us define $C_0 = U$, $C_{n+1} := \emptyset$ and, for $1 \leq k \leq n$,

$$C_k := \bigcup_{i_1 < \dots < i_k} (A_{i_1} \cap A_{i_2} \cdots \cap A_{i_k}).$$

Then for $1 \leq k \leq n$, C_k is the union of all possible intersections of the k -subsets of the collection Γ . In particular $C_1 = A_0 \cup A_1 \cup \dots \cup A_{n-1}$ and $C_n = A_0 \cap A_1 \cap \dots \cap A_{n-1}$. Let $B_k := C_k - C_{k+1}$ for $0 \leq k \leq n$. Observe that $C_k \supseteq C_{k+1}$, so $|B_k| = |C_k| - |C_{k+1}|$. Let us also define

$$b_k := |B_k| \text{ and } t_k := \sum_{i_1 < \dots < i_k} |A_{i_1} \cap A_{i_2} \cdots \cap A_{i_k}|.$$

We have $t_1 = \sum_{i=0}^{n-1} |A_i|$ and $t_2 = \sum_{0 \leq i < j < n} |A_i \cap A_j|$. We also have $b_n = t_n$ and $b_{n-1} = t_{n-1} - \binom{n}{1} b_n$. Now $A_0 \cap A_1 \cap \dots \cap A_{n-1}$ is a subset of $\binom{n}{n-2} = \binom{n}{2}$ sets of the form $A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_{n-2}}$ and each $(n-1)$ -section $A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_{n-1}}$ is a subset of $\binom{n-1}{n-2} = \binom{n-1}{1}$ sets of the form $A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_{n-2}}$ with $i_1 < i_2 < \dots < i_{n-2}$. Therefore $b_{n-2} = t_{n-2} - \binom{n-1}{1} b_{n-1} - \binom{n}{2} b_n$.

Similarly, since each $(n-k+i)$ -section is a subset of $\binom{n-k+i}{i}$ intersections of $(n-k)$ sets for $i = 1, \dots, k$, we have the recursive formula

$$b_{n-k} = t_{n-k} - \sum_{i=1}^k \binom{n-k+i}{i} b_{n-k+i} \text{ for } k = 0, \dots, n. \quad (1)$$

Lemma 2 *Following the notation introduced above*

$$b_{n-k} = \sum_{i=0}^k (-1)^i \binom{n-k+i}{i} t_{n-k+i} \text{ for } k = 0, \dots, n. \quad (2)$$

Proof: We will prove the claim by strong induction on k . For $k = 0$ we have $b_n = t_n$. Now assume that the claim is true for any integer $i \leq k$ in place of k .

By the recursive formula (1) we have

$$b_{n-(k+1)} = t_{n-(k+1)} - \binom{n-k}{1} b_{n-k} - \binom{n-k+1}{2} b_{n-k+1} - \dots - \binom{n}{k+1} b_n.$$

On the other hand by induction we have the following equations

$$\begin{aligned}
b_n &= t_n \\
b_{n-1} &= t_{n-1} - \binom{n}{1} t_n \\
b_{n-2} &= t_{n-2} - \binom{n-1}{1} t_{n-1} + \binom{n}{2} t_n \\
&\vdots \\
b_{n-k} &= t_{n-k} - \binom{n-k+1}{1} t_{n-k+1} + \cdots + (-1)^k \binom{n}{k} t_n.
\end{aligned}$$

It follows that

$$\begin{aligned}
-\binom{n}{k+1} b_n &= -\binom{n}{k+1} t_n \\
-\binom{n-1}{k} b_{n-1} &= -\binom{n-1}{k} t_{n-1} + \binom{n-1}{k} \binom{n}{1} t_n \\
-\binom{n-2}{k-1} b_{n-2} &= -\binom{n-2}{k-1} t_{n-2} + \binom{n-2}{k-1} \binom{n-1}{1} t_{n-1} - \binom{n-2}{k-1} \binom{n}{2} t_n \\
&\vdots \\
-\binom{n-k}{1} b_{n-k} &= -\binom{n-k}{1} t_{n-k} + \binom{n-k}{1} \binom{n-k+1}{1} t_{n-k+1} - \cdots - (-1)^{k+1} \binom{n-k}{1} \binom{n}{k} t_n.
\end{aligned}$$

If we sum all these equalities, then on the right hand side the coefficient of t_n is

$$c(t_n) = \sum_{i=0}^k (-1)^{k-i+1} \binom{n-k+i}{i+1} \binom{n}{k-i}. \text{ For } d \leq k \text{ one has}$$

$$\binom{n-d}{k-d+1} \binom{n}{d} = \frac{(n-d)!}{(n-k-1)!(k-d+1)!} \frac{n!}{(n-d)!d!} = \frac{n!}{(k+1)!(n-k-1)!} \frac{(k+1)!}{d!(k-d+1)!} = \binom{n}{k+1} \binom{k+1}{d}.$$

$$\text{Then } c(t_n) = \binom{n}{k+1} \sum_{i=0}^k (-1)^{k-i+1} \binom{k+1}{k-i} = -\binom{n}{k+1} (-1)^k = (-1)^{k+1} \binom{n}{k+1},$$

where the last equality follows from the fact that

$$\binom{k+1}{0} - \binom{k+1}{1} + \binom{k+1}{2} + \cdots + (-1)^k \binom{k+1}{k} = -(-1)^{k+1} = (-1)^k.$$

$$\text{Similarly for } s = 1, \dots, k \text{ we have } c(t_{n-s}) = \sum_{i=0}^{k-s} (-1)^{k-s-i+1} \binom{n-s-k+i}{i+1} \binom{n}{k-s-i}$$

$$= \binom{n-s}{k-s+1} \sum_{i=0}^{k-s} (-1)^{k-s-i+1} \binom{n}{k-s-i} = (-1)^{k-s+1} \binom{n-s}{k-s+1}.$$

Now plugging $s = k - i$ in the formula above we get

$$b_{n-(k+1)} = \sum_{i=0}^k (-1)^{i+1} \binom{n-k+i}{i+1} t_{n-k+i}. \quad \square$$

Proposition 1 *Following the same notation one has for $1 \leq k \leq n$,*

$$\sum_{i=0}^n i^k b_i = \sum_{i=1}^k i^k \left[\sum_{j=i}^k (-1)^{j-i} \binom{j}{j-i} t_j \right], \text{ In particular:}$$

$$(a) \sum_{i=0}^n i b_i = t_1 = \sum_{i=0}^{n-1} |A_i| \text{ (Inclusion Exclusion Principle) and}$$

$$(b) \sum_{i=0}^n i^2 b_i = t_1 + 2t_2 = \sum_{i=0}^{n-1} |A_i| + 2 \sum_{i < j} |A_i \cap A_j|.$$

Proof: According to Lemma 2 we have

$$\begin{aligned}
b_n &= t_n \\
b_{n-1} &= t_{n-1} - \binom{n}{1}t_n \\
b_{n-2} &= t_{n-2} - \binom{n-1}{1}t_{n-1} + \binom{n}{2}t_n \\
&\vdots \\
b_2 &= t_2 - \binom{3}{1}t_3 + \binom{4}{2}t_4 + \cdots + (-1)^{n-3}\binom{n-1}{n-3}t_{n-1} + (-1)^{n-2}\binom{n}{n-2}t_n \\
b_1 &= t_1 - \binom{2}{1}t_2 + \binom{3}{2}t_3 + \binom{4}{3}t_4 + \cdots + (-1)^{n-2}\binom{n-1}{n-2}t_{n-1} + (-1)^{n-1}\binom{n}{n-1}t_n
\end{aligned}$$

If we sum $\sum_{i=1}^n i^k b_i$, then for $1 \leq s \leq n$, the coefficient of t_s on the right hand side is

$$c(t_s) = \sum_{i=1}^s i^k \binom{s}{s-i} (-1)^{s-i}.$$

We claim that $c(t_s) = 0$ for $k < s \leq n$. We prove this by strong induction on k . For $k = 1$ we have

$$c(t_s) = \sum_{i=1}^s i \binom{s}{s-i} (-1)^{s-i} = \sum_{i=1}^s i \binom{s}{i} (-1)^{s-i} = \sum_{i=1}^s s \binom{s-1}{i-1} (-1)^{s-i}$$

Since $s \geq 2$, by substituting $m = s - 1 \geq 1$ and $j = i - 1$

$$c(t_s) = s \sum_{j=0}^m \binom{m}{j} (-1)^{m-j} = s(1 - 1)^m = 0.$$

Now assume $\sum_{i=1}^s i^l \binom{s}{s-i} (-1)^{s-i} = 0$ for any $1 \leq l \leq k$ and $l + 1 \leq s \leq n$. Then

$$c(t_s) = \sum_{i=1}^s i^{k+1} \binom{s}{s-i} (-1)^{s-i} = \sum_{i=1}^s i^{k+1} \binom{s}{i} (-1)^{s-i} = \sum_{i=1}^s s i^k \binom{s-1}{i-1} (-1)^{s-i}.$$

Substituting $m = s - 1 \geq l \geq 1$ and $j = i - 1$ we obtain

$$\begin{aligned}
c(t_s) &= s \sum_{j=0}^m (j+1)^k \binom{m}{j} (-1)^{m-j} = s \sum_{j=0}^m \sum_{l=0}^k \binom{k}{l} j^l \binom{m}{j} (-1)^{m-j} \\
&= s \sum_{l=0}^k \binom{k}{l} \sum_{j=0}^m j^l \binom{m}{j} (-1)^{m-j}.
\end{aligned}$$

Since $m = s - 1 \geq l$, we have $m \geq l + 1$ and by induction hypothesis each summand

$$\sum_{j=0}^m j^l \binom{m}{j} (-1)^{m-j} = \sum_{j=1}^m j^l \binom{m}{m-j} (-1)^{m-j} = 0.$$

Hence $c(t_s) = 0$. On the other hand for $1 \leq s \leq k$ the coefficient of each i^s on the right hand side is $\sum_{j=i}^k (-1)^{j-i} \binom{j}{j-i} t_j$. Hence we have the result. In particular, for $k = 2$ the non-zero terms on the right-hand-side are $t_1 - \binom{2}{1}t_2 + 2^2t_2 = t_1 + 2t_2$. \square

Theorem 1 Let $\phi(n) = |\{1 \leq i \leq n : \gcd(i, n) = 1\}|$ denote Euler's totient function. Let X be a random variable which counts the number of distinct roots of a monic polynomial in $\mathbb{Z}_n[x]$ of degree $m > 0$. Then

(a) $E[X] = 1$ and

(b) if $m = 1$ then $\text{Var}[X] = 0$, otherwise $\text{Var}[X] = \sum_{d|n, d \neq n} \frac{d}{n} \phi(\frac{n}{d}) = \sum_{d|n} \frac{d-1}{n} \phi(\frac{n}{d})$.
In particular, if $n = p^k$ where p is a prime number and $k \geq 1$, $\text{Var}[X] = k(1 - 1/p)$.

Remark 1. We found this result by direct computation and using the Online Encyclopedia of Integer Sequences (OEIS) see [7]. For polynomials of degree 2,3,4,5 in $\mathbb{Z}_n[x]$ we computed $E[X]$ and $\text{Var}[X]$ for $n = 2, 3, 4, \dots, 20$ using Maple and found that $E[X] = 1$ in all cases. Values for the variance are given in the table below.

n	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$\text{Var}[X]$	$\frac{1}{2}$	$\frac{2}{3}$	1	$\frac{4}{5}$	$\frac{3}{2}$	$\frac{6}{7}$	$\frac{3}{2}$	$\frac{4}{3}$	$\frac{17}{10}$	$\frac{10}{11}$	$\frac{7}{3}$	$\frac{12}{13}$	$\frac{25}{14}$	2	2
$a(n)$	1	2	4	4	9	6	12	12	17	10	28	12	25	30	32

When we first computed $\text{Var}[X]$ we did not recognize the numbers. Writing $\text{Var}[X] = a(n)/n$ we computed the sequence for $a(n)$ (see the table) and looked it up in the OEIS. We found it is sequence A006579 and that $a(n) = \sum_{k=1}^{n-1} \gcd(n, k)$. The OEIS also has the formula $a(n) = \sum_{d|n} (d-1)\phi(\frac{n}{d})$.

Proof: Let A_i be the set of all monic univariate polynomials of degree $m > 0$ which have a root at $\alpha_i \in \mathbb{Z}_n$. Then since $x - \alpha_i$ is monic, for any $f \in A_i$ we have $f = (x - \alpha_i)q$ for a unique $q \in \mathbb{Z}_n[x]$ and we have n^{m-1} choices for such an f . Hence $|A_i| = n^{m-1}$.

Let $x_i := \text{Prob}[X = i]$. This is the probability that f has exactly i distinct roots, i.e. $f \in B_i$ in the notation introduced in section 1 considering the finite collection of sets $\Gamma = \{A_i, i = 0, \dots, n-1\}$. Since we have n^{m-1} choices for a monic polynomial of degree m in $\mathbb{Z}_n[x]$ we have $x_i = \frac{b_i}{n^{m-1}}$. Then by Proposition 1

$$E[X] = \sum_{i=0}^n i x_i = \sum_{i=0}^n i \frac{b_i}{n^{m-1}} = \frac{\sum_{i=0}^n i b_i}{n^{m-1}} = \frac{\sum_{i=0}^{n-1} |A_i|}{n^{m-1}} = \frac{\sum_{i=0}^{n-1} n^{m-1}}{n^{m-1}} = \frac{nn^{m-1}}{n^{m-1}} = 1$$

To prove (b), if $m = 1$ then $f = x - \alpha$ for some $\alpha \in \mathbb{Z}_n$ and hence $X = 1$ and $\text{Var}[X] = 0$. For $m > 1$ and $\alpha \in \mathbb{Z}_n^*$, our first aim is to find $|A_0 \cap A_\alpha|$. Let $f \in A_0 \cap A_\alpha$. It may not be the case that $f = x(x - \alpha)q$ for a unique $q \in \mathbb{Z}_n[x]$, since $\mathbb{Z}_n[x]$ is not a unique factorization domain in general. However $f = xq_1 = (x - \alpha)q_2$ for unique $q_1, q_2 \in \mathbb{Z}_n[x]$. It follows that $\alpha q_2(0) = 0 \pmod n$. If $\gcd(\alpha, n) = d$ then $\gcd(\frac{\alpha}{d}, \frac{n}{d}) = 1$ and hence $q_2(0) = 0 \pmod{\frac{n}{d}}$. The general form of $q_2 = x^{m-1} + a_{m-2}x^{m-2} + \dots + a_0$ where $a_i \in \mathbb{Z}_n$ for $i = 0, \dots, m-2$. Since $q_2(0) = a_0 \pmod{\frac{n}{d}}$, there are d choices for a_0 and hence there are dn^{m-2} choices for q_2 . Therefore $|A_0 \cap A_\alpha| = dn^{m-2}$.

For a given pair (γ, β) with $\beta > \gamma$, to compute $|A_\gamma \cap A_\beta|$, define $\alpha := \beta - \gamma$ and consider $A_0 \cap A_\alpha$. If $f \in A_\gamma \cap A_\beta$, then we have $f(x) = (x - \gamma)q_3(x) = (x - \beta)q_4(x)$ for unique $q_3, q_4 \in \mathbb{Z}_n[x]$. By the coordinate translation $x \mapsto x + \gamma$ we have $f(x + \gamma) \in A_0 \cap A_\alpha$, since $f(x + \gamma) = xq_3(x + \gamma) = (x - \alpha)q_4(x + \gamma)$ where $f(x + \gamma), q_3(x + \gamma), q_4(x + \gamma)$ are monic and with the same degree before the translation. This correspondence is bijective and it follows that $|A_\gamma \cap A_\beta| = |A_0 \cap A_\alpha| = dn^{m-2}$.

Let $d = \gcd(\alpha, n)$. There are $k = \phi(\frac{n}{d})$ elements β_1, \dots, β_k in $\mathbb{Z}_{\frac{n}{d}}$ such that $\gcd(\beta_j, \frac{n}{d}) = 1$. If we define $\alpha_j := d\beta_j \in \mathbb{Z}_n$ then $\gcd(\alpha_j, n) = d$. For, if $s = \gcd(\alpha_j, n)$ and $d|s$ then $s|\alpha_j \Rightarrow s|d\beta_j \Rightarrow \frac{s}{d}|\beta_j$

and $\frac{s}{d} | \frac{n}{d} \Rightarrow \frac{s}{d} | \gcd(\beta_j, \frac{n}{d}) \Rightarrow \frac{s}{d} | 1 \Rightarrow s = d$. Now, for each j consider the $n - \alpha_j$ pairs of the form $(i, i + \alpha_j)$ where $i = 0, \dots, n - \alpha_j - 1$. We have $|A_i \cap A_{i+\alpha_j}| = |A_0 \cap A_{\alpha_j}|$ and

$$\sum_{\beta > \gamma, d = \gcd(\beta - \gamma, n)} |A_\gamma \cap A_\beta| = \sum_{j=1}^k (n - \alpha_j) |A_0 \cap A_{\alpha_j}| = \sum_{j=1}^k (n - \alpha_j) d n^{m-2} = d n^{m-2} \sum_{j=1}^k n - \alpha_j$$

where $d = \gcd(\alpha_j, n)$ and $k = \phi(\frac{n}{d})$. Since $\gcd(n, \alpha_j) = d \iff \gcd(n, n - \alpha_j) = d$ we have $\sum_{j=1}^k n - \alpha_j = \sum_{j=1}^k \alpha_j$. Then

$$2 \sum_{j=1}^k \alpha_j = \sum_{j=1}^k \alpha_j + \sum_{j=1}^k n - \alpha_j = \sum_{j=1}^k n = k n = \phi(\frac{n}{d}) n \implies \sum_{j=1}^k \alpha_j = \frac{n}{2} \phi(\frac{n}{d}).$$

It follows that

$$\sum_{\beta > \gamma, d = \gcd(\beta - \gamma, n)} |A_\gamma \cap A_\beta| = d n^{m-2} \sum_{j=1}^k n - \alpha_j = d n^{m-2} \sum_{j=1}^k \alpha_j = \frac{n}{2} \phi(\frac{n}{d}) d n^{m-2}.$$

Then by Proposition 1 it follows that

$$\begin{aligned} \text{Var}[X] &= \text{E}[X^2] - \text{E}[X]^2 = -1^2 + \text{E}[X^2] \\ &= -1 + \sum_{i=0}^n i^2 x_i = -1 + \sum_{i=0}^n i^2 \frac{b_i}{n^m} = -1 + \frac{\sum_{i=0}^n i^2 b_i}{n^m} \\ &= -1 + \frac{\sum_{i=0}^{n-1} |A_i| + 2 \sum_{i < j} |A_i \cap A_j|}{n^m} \\ &= -1 + \frac{n n^{m-1}}{n^m} + \frac{2 \sum_{d|n, d \neq n} \frac{n}{2} \phi(\frac{n}{d}) d n^{m-2}}{n^m} \\ &= 2 \sum_{d|n, d \neq n} \frac{n}{2} \phi(\frac{n}{d}) d n^{-2} = \sum_{d|n, d \neq n} \frac{d}{n} \phi(\frac{n}{d}). \end{aligned}$$

Also, since by Gauss' Lemma $\sum_{d|n} \phi(\frac{n}{d}) = n$ we have

$$\begin{aligned} \sum_{d|n} \frac{d-1}{n} \phi(\frac{n}{d}) &= \sum_{d|n} \frac{d}{n} \phi(\frac{n}{d}) - \frac{1}{n} \sum_{d|n} \phi(\frac{n}{d}) \\ &= \phi(1) + \sum_{d|n, d \neq n} \frac{d}{n} \phi(\frac{n}{d}) - \frac{1}{n} n = \sum_{d|n, d \neq n} \frac{d}{n} \phi(\frac{n}{d}). \end{aligned}$$

To prove the last claim, let $n = p^k$ where p is a prime number and $k \geq 1$. Then

$$\sum_{d|n, d \neq n} \frac{d}{n} \phi(\frac{n}{d}) = \sum_{s=0}^{k-1} \frac{p^s}{p^k} \phi(\frac{p^k}{p^s}) = \sum_{s=0}^{k-1} p^{s-k} p^{k-s-1} (p-1) = k(1 - 1/p).$$

□

Theorem 2 Let f, g be polynomials in $\mathbb{F}_q[x, y]$ of the form $f = c_n x^n + \sum_{i=0}^{n-1} \sum_{j=0}^{n-i} c_{ij} x^i y^j$ and $g = d_m x^m + \sum_{i=0}^{m-1} \sum_{j=0}^{m-i} d_{ij} x^i y^j$ with $c_n \neq 0$ and $d_m \neq 0$, thus of total degree n and m respectively. Let X be a random variable that counts the number of $\gamma \in \mathbb{F}_q$ such that $\gcd(f(x, \gamma), g(x, \gamma)) \neq 1$. If $n > 0$ and $m > 0$ then

- (a) $E[X] = 1$ and
- (b) $\text{Var}[X] = 1 - 1/q$.

Remark 2. We found this result by computation. For quadratic polynomials f, g of the form $f = x^2 + (a_1 y + a_2)x + a_3 y^2 + a_4 y + a_5$ and $g = x^2 + (b_1 y + b_2)x + b_3 y^2 + b_4 y + b_5$ over finite fields of size $q = 2, 3, 4, 5, 8, 9, 11$ we generated all q^{10} pairs and computed $X = |\{\alpha \in \mathbb{F}_q : \gcd(f(x, \alpha), g(x, \alpha)) \neq 1\}|$. Magma code for \mathbb{F}_4 is given in Appendix A. We repeated this for cubic polynomials and some higher degree bivariate polynomials for $q = 2, 3$ to verify that $E[X] = 1$ and $\text{Var}[X] = 1 - 1/q$ holds more generally. For yet higher degree polynomials we used random samples. That $E[X] = 1$ independent of the degrees of f and g was a surprise to us. We had expected a logarithmic dependence on the degrees of f and g .

Proof: Without loss of generality we may assume f and g are monic in x because $\gcd(f(x, \gamma), g(x, \gamma)) = 1 \iff \gcd(c_n^{-1}f(x, \gamma), d_m^{-1}g(x, \gamma)) = 1$. For $\gamma \in \mathbb{F}_q$, let us define A_γ as the set of polynomial pairs $(f, g) \in \mathbb{F}_q[x, y]^2$ where f, g are monic in x with total degrees, $\deg(f) = n > 0$ and $\deg(g) = m > 0$ such that $\gcd(f(x, \gamma), g(x, \gamma)) \neq 1$. Our first aim is to compute $|A_0|$.

Let $(f, g) \in A_0$. Since f and g are monic in x , $f(x, 0), g(x, 0)$ are monic polynomials of degree n and m respectively in $\mathbb{F}_q[x]$. We have finitely many choices, say s , for non-relatively prime monic polynomial pairs $(h_i(x), l_i(x))$ with $\deg(h_i) = n$ and $\deg(l_i) = m$ with $i = 1, \dots, s$ in $\mathbb{F}_q[x]^2$. Let $(f(x, 0), g(x, 0)) = (h_i(x), l_i(x))$ for some fixed i where $1 \leq i \leq s$. In fact $s = (q^n q^m)/q = q^{n+m-1}$, since there are $q^n q^m$ possible choices for monic polynomial pairs (h, l) in $\mathbb{F}_q[x]$ with $\deg(h) = n$, $\deg(l) = m$ and the probability of a given monic pair is non-relatively prime over $\mathbb{F}_q[x]$ is $1/q$ (see [8, 2] and also [1] for an accessible proof).

Let $f(x, y) = x^n + c_{n-1}(y)x^{n-1} + \dots + c_1(y)x + c_0(y)$ where $c_d(y) \in \mathbb{F}_q[y]$ of total degree $\deg(c_{n-d}(y)) \leq d$ and let $c_{n-d}(y) = a_d^{(n-d)}y^d + \dots + a_0^{(n-d)}$ where $a_i^{(n-d)} \in \mathbb{F}_q$.

Let $h_i(x) = x^n + \alpha_{n-1}^{(i)}x^{n-1} + \dots + \alpha_0^{(i)}$ with $\alpha_v^{(i)} \in \mathbb{F}_q$ for $0 \leq v \leq n-1$. Then for $1 \leq d \leq n$, we have $c_{n-d}(0) = a_0^{(n-d)} = \alpha_{n-d}^{(i)}$. It follows that there are q^d choices for such $c_{n-d}(y)$ and hence there are $q^1 q^2 \dots q^n = q^{n(n+1)/2}$ choices for such $f(x, y)$. Similarly there are $q^{m(m+1)/2}$ choices for $g(x, y)$. Let us denote these numbers as $D = q^{n(n+1)/2}$ and $R = q^{m(m+1)/2}$. Since we have s choices for i , $|A_0| = sDR$.

On the other hand for a given $\gamma \in \mathbb{F}_q$ if $(f(x, y), g(x, y)) \in A_0$ then $(f(x, y - \gamma), g(x, y - \gamma)) \in A_\gamma$, since $f(x, y - \gamma)$ is again a bivariate polynomial which is a monic polynomial in x of total degree n and $g(x, y - \gamma)$ is again a bivariate polynomial which is monic polynomial in x of total degree m . This correspondence (coordinate transformation) is bijective. Hence for any $\gamma \in \mathbb{F}_q$, one has $|A_\gamma| = sDR$.

For a general polynomial $f(x, y) \in \mathbb{F}_q[x, y]$ which is monic in x and of total degree $n > 0$, one has $q^2 q^3 \dots q^{n+1} = q^n D$ choices. Similarly for a general polynomial $g(x, y) \in \mathbb{F}_q[x, y]$ which is monic in x and of total degree $m > 0$, one has $q^2 q^3 \dots q^{m+1} = q^m R$ choices and therefore there are $q^{n+m} DR$ pairs (f, g) which are monic in x with total degrees $\deg(f) = n$ and $\deg(g) = m$.

Let $x_i := \text{Prob}[X = i]$. This is the probability that $\gcd(f(x, \gamma), g(x, \gamma)) \neq 1$ for exactly i different γ 's in \mathbb{F}_q , i.e. the probability that $(f, g) \in B_i$ in the notation introduced in section 1 considering the finite

collection of sets $\Gamma = \{A_\gamma, \gamma \in \mathbb{F}_q\}$. Hence $x_i = \frac{b_i}{q^{n+m}DR}$. Then by Proposition 1

$$\begin{aligned} \mathbb{E}[X] &= \sum_{i=0}^q ix_i = \sum_{i=0}^q i \frac{b_i}{q^{n+m}DR} = \frac{\sum_{i=0}^q ib_i}{q^{n+m}DR} \\ &= \frac{\sum_{i=0}^{q-1} |A_i|}{q^{n+m}DR} = \frac{\sum_{i=0}^{q-1} sDR}{q^{n+m}DR} = \frac{qsDR}{q^{n+m}DR} = \frac{qq^{n+m-1}}{q^{n+m}} = 1. \end{aligned}$$

To determine the variance of X , our proof assumes a set ordering of the elements of \mathbb{F}_q . For this purpose let us fix a generator α of \mathbb{F}_q^* and use the ordering $0 < 1 < \alpha < \alpha^2 < \dots < \alpha^{q-2}$.

For $(\gamma, \theta) \in \mathbb{F}_q^2$ with $\gamma < \theta$, let us define $A_{\gamma, \theta}$ as the set of bivariate polynomial pairs (f, g) with f, g are monic in x with total degrees, $\deg(f) = n > 0$ and $\deg(g) = m > 0$ such that $\gcd(f(x, \gamma), g(x, \gamma)) \neq 1$ and $\gcd(f(x, \theta), g(x, \theta)) \neq 1$. Our first aim is to compute $|A_{0,1}|$.

Let $f, g \in A_{0,1}$. Since f and g are monic in x , $f(x, 0), f(x, 1)$ are monic polynomials of degree n and $g(x, 0), g(x, 1)$ are monic polynomials of degree m in $\mathbb{F}_q[x]$. We have finitely many choices for non-relatively prime monic polynomial pairs $(h_i(x), l_i(x))$ with $\deg(h_i) = n$ and $\deg(l_i) = m$ with $i = 1, \dots, s$ in $\mathbb{F}_q[x]^2$.

Let $(f(x, 0), g(x, 0)) = (h_i(x), l_i(x))$ and $(f(x, 1), g(x, 1)) = (h_j(x), l_j(x))$ for some fixed pair (i, j) where $1 \leq i, j \leq s$.

Let $f(x, y) = x^n + c_{n-1}(y)x^{n-1} + \dots + c_1(y)x + c_0(y)$ where $c_d(y) \in \mathbb{F}_q[y]$ of total degree $\deg(c_{n-d}(y)) \leq d$ and let $c_{n-d}(y) = a_d^{(n-d)}y^d + \dots + a_0^{(n-d)}$ where $a_i^{(n-d)} \in \mathbb{F}_q$.

Let $h_i(x) = x^n + \alpha_{n-1}^{(i)}x^{n-1} + \dots + \alpha_0^{(i)}$ and $h_j(x) = x^n + \beta_{n-1}^{(j)}x^{n-1} + \dots + \beta_0^{(j)}$ with $\alpha_v^{(i)}, \beta_w^{(j)} \in \mathbb{F}_q$ for $0 \leq v, w \leq n-1$. Then for $1 \leq d \leq n$, we have

$$c_{n-d}(0) = a_0^{(n-d)} = \alpha_{n-d}^{(i)} \quad \text{and} \quad c_{n-d}(1) = a_d^{(n-d)} + \dots + a_1^{(n-d)} + a_0^{(n-d)} = \beta_{n-d}^{(j)}.$$

It follows that there are q^{d-1} choices for such $c_{n-d}(y)$ and hence there are $q^0 q^1 \dots q^{n-1} = q^{n(n-1)/2}$ choices for such $f(x, y)$. Similarly there are $q^{m(m-1)/2}$ choices for $g(x, y)$. Let us call these numbers as $D_1 = q^{n(n-1)/2}$ and $R_1 = q^{m(m-1)/2}$. Since we have s^2 choices for (i, j) (i and j are need not be different, $|A_{0,1}| = s^2 D_1 R_1$).

On the other hand if $(f(x, y), g(x, y)) \in A_{0,1}$ then for $\gamma, \theta \in \mathbb{F}_q$ with $\gamma < \theta$, $(f(x, \frac{y-\gamma}{\theta-\gamma}), g(x, \frac{y-\gamma}{\theta-\gamma})) \in A_{\gamma, \theta}$, since $f(x, \frac{y-\gamma}{\theta-\gamma})$ is again a monic polynomial in x of total degree n and $g(x, \frac{y-\gamma}{\theta-\gamma})$ is again a monic polynomial in x of total degree m . This correspondence (coordinate transformation) is bijective and preserves relative primeness. Hence for a given $\gamma, \theta \in \mathbb{F}_q$ with $\gamma < \theta$, one has $|A_{\gamma, \theta}| = s^2 D_1 R_1$.

For a general bivariate polynomial $f(x, y) \in \mathbb{F}_q[x, y]$ which is monic in x and of total degree n , one has $q^2 q^3 \dots q^{n+1} = q^{2n} D_1$ choices. Similarly for a general bivariate polynomial $g(x, y) \in \mathbb{F}_q[x, y]$ which is monic in x and of total degree m , one has $q^2 q^3 \dots q^{m+1} = q^{2m} R_1$ choices and therefore the number of bivariate polynomial pairs in (f, g) which are monic in x with total degrees, $\deg(f) = n$ and $\deg(g) = m$ is $q^{2n+2m} D_1 R_1$. Then with this notation we have $x_i = \frac{b_i}{q^{2n+2m} D_1 R_1}$.

Since we have $\binom{q}{2}$ choices for (γ, θ) with $\gamma < \theta$, $|A_{\gamma, \theta}| = s^2 D_1 R_1$ for all (γ, θ) with $\gamma < \theta$ and $\mathbb{E}[X] = 1$, by Proposition 1 we have

$$\begin{aligned}
\text{Var}[X] &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \mathbb{E}[X^2] - 1^2 = -1 + \sum_{i=0}^q i^2 x_i \\
&= -1 + \sum_{i=0}^q i^2 \frac{b_i}{q^{2n+2m} D_1 R_1} = -1 + \frac{\sum_{i=0}^q i^2 b_i}{q^{2n+2m} D_1 R_1} \\
&= -1 + \frac{\sum_{i=0}^{q-1} |A_i| + 2 \sum_{i<j} |A_i \cap A_j|}{q^{2n+2m} D_1 R_1} = -1 + \frac{\sum_{i=0}^{q-1} |A_i|}{q^{2n+2m} D_1 R_1} + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} \\
&= -1 + \frac{\sum_{i=0}^q i b_i}{q^{2n+2m} D_1 R_1} + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} = -1 + \sum_{i=0}^q i \frac{b_i}{q^{2n+2m} D_1 R_1} + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} \\
&= -1 + \sum_{i=0}^q i x_i + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} = -1 + \mathbb{E}[X] + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} \\
&= -1 + 1 + \frac{2 \sum_{i<j} s^2 D_1 R_1}{q^{2n+2m} D_1 R_1} = \frac{2 \binom{q}{2} s^2 D_1 R_1}{q^{2n} q^{2m} D_1 R_1} = \frac{q(q-1)q^{2n+2m-2}}{q^{2n+2m}} \\
&= \frac{q(q-1)}{q^2} = 1 - \frac{1}{q}. \quad \square
\end{aligned}$$

Theorem 3 Let $f, g \in \mathbb{F}_q[x_1, x_2, \dots, x_n]$ be of the form $f = c_l x_1^l + \sum_{i=0}^{l-1} c_{l-i}(x_2, \dots, x_n)x^i$ and $g = d_m x_1^m + \sum_{i=0}^{m-1} d_{m-i}(x_2, \dots, x_n)x^i$ where $c_l \neq 0, d_m \neq 0, \deg c_{l-i} \leq l-i$, and $\deg d_{m-i} \leq m-i$, thus f and g have total degree l and m respectively. Let X be a random variable which counts the number of $\gamma = (\gamma_2, \dots, \gamma_n) \in \mathbb{F}_q^{n-1}$ such that $\gcd(f(x_1, \gamma_2, \dots, \gamma_n), g(x_1, \gamma_2, \dots, \gamma_n)) \neq 1$. If $n > 1, l > 0$ and $m > 0$ then

- (a) $\mathbb{E}[X] = q^{n-2}$ and
- (b) $\text{Var}[X] = q^{n-2}(1 - 1/q)$.

It follows from (a) that if γ is chosen at random from \mathbb{F}_q^{n-1} then

$$\text{Prob}[\gcd(f(x_1, \gamma_2, \dots, \gamma_n), g(x_2, \gamma_2, \dots, \gamma_n)) \neq 1] = \frac{q^{n-2}}{q^{n-1}} = \frac{1}{q}.$$

Proof: The proof runs along the same lines of the proof of Theorem 2. Let U be the set of all possible monic pairs $(f, g) \in \mathbb{F}_q[x_1, \dots, x_n]^2$ where f, g are as described in the corollary and for $\alpha = (\alpha_2, \dots, \alpha_n) \in \mathbb{F}_q^{n-1}$, let A_α be the set of all such polynomial pairs with the property that $\gcd(f(x_1, \alpha), g(x_1, \alpha)) \neq 1$.

For the first part we will consider the monic pairs (f, g) with

$$\gcd(f(x_1, 0, 0, \dots, 0), g(x_1, 0, 0, \dots, 0)) \neq 1$$

and compute that probability of this event is $1/q$ again. Then for a given non-zero $\alpha = (\alpha_2, \dots, \alpha_n) \in \mathbb{F}_q^{n-1}$ considering the coordinate change

$$\bar{f}(x_1, \alpha_2, \dots, \alpha_n) = f(x_1, x_2 - \alpha_2, \dots, x_n - \alpha_n)$$

and using Proposition 1 part (a), since there are q^{n-1} possible such α 's, we will see that $E[X] = q^{n-1}q^{-1} = q^{n-2}$.

For the second part we will consider the monic pairs (f, g) with

$$\begin{aligned} \gcd(f(x_1, 0, 0 \dots, 0), g(x_1, 0, 0 \dots, 0)) &\neq 1 \\ \gcd(f(x_1, 1, 0 \dots, 0), g(x_1, 1, 0 \dots, 0)) &\neq 1 \end{aligned}$$

and see that probability of this event is $1/q^2$ again. For a given pair $(\alpha, \beta) \in \mathbb{F}_q^{n-1} \times \mathbb{F}_q^{n-1}$ with $\alpha \neq \beta$, this time the coordinate change of the second part of the proof that computes the variance may not be that obvious. We give the explicit construction below. Then by enumerating the elements of \mathbb{F}_q^{n-1} from 0 to $q^{n-1} - 1$ and using Proposition 1 part (b), since there are $\binom{q^{n-1}}{2}$ possible pairs (α, β) with $\alpha < \beta$, we will see that

$$\begin{aligned} \text{Var}[X] &= E[X^2] - E[X]^2 = -E[X]^2 + \sum_{i=0}^{q^{n-1}-1} i^2 \Pr(X = i) \\ &= -E[X]^2 + \sum_{i=0}^{q^{n-1}-1} i^2 \frac{b_i}{|U|} = -E[X]^2 + \frac{\sum_{i=0}^{q^{n-1}-1} i^2 b_i}{|U|} \\ &= -E[X]^2 + \frac{\sum_{i=0}^{q^{n-1}-1} |A_i| + 2 \sum_{i < j} |A_i \cap A_j|}{|U|} \\ &= -E[X]^2 + \sum_{i=0}^{q^{n-1}-1} \frac{1}{q} + 2 \sum_{i < j} \frac{1}{q^2} \\ &= -E[X]^2 + q^{n-1} \frac{1}{q} + 2 \binom{q^{n-1}}{2} \frac{1}{q^2} \\ &= -q^{2n-4} + q^{n-2} + q^{n-1} (q^{n-1} - 1) q^{-2} \\ &= q^{n-2} - q^{n-3} = q^{n-2} \left(1 - \frac{1}{q}\right). \end{aligned}$$

For a given pair $(\alpha, \beta) \in \mathbb{F}_p^{n-1} \times \mathbb{F}_p^{n-1}$ with $\alpha \neq \beta$, let $\alpha = (\alpha_2, \dots, \alpha_n)$ and $\beta = (\beta_2, \dots, \beta_n)$. Our aim is to find a coordinate change such that

$$\begin{aligned} \bar{f}(x_1, \alpha_2, \dots, \alpha_n) &= f(x_1, 0, 0 \dots, 0) \\ \bar{f}(x_1, \beta_2, \dots, \beta_n) &= f(x_1, 1, 0 \dots, 0) \end{aligned}$$

where

$$\bar{f}(x_1, x_2, \dots, x_n) = f(x_1, a_{20} + a_{22}x_2 + \dots + a_{2n}x_n, \dots, a_{n0} + a_{n2}x_2 + \dots + a_{nn}x_n).$$

Note that this transformation does not change the leading term in x_1 , so it preserves monicness and degree in x_1 and preserves coprimality. To make this transformation bijective we need a $(n-1) \times (n-1)$ matrix

$$A = \begin{pmatrix} a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots \\ a_{n2} & \dots & a_{nn} \end{pmatrix}$$

which is invertible and to satisfy the relations we need

$$\begin{aligned} a_{22}\alpha_2 + \cdots + a_{2n}\alpha_n &= -a_{20} \\ a_{22}\beta_2 + \cdots + a_{2n}\beta_n &= 1 - a_{20} \end{aligned}$$

and for $3 \leq j \leq n$

$$\begin{aligned} a_{j2}\alpha_2 + \cdots + a_{jn}\alpha_n &= -a_{j0} \\ a_{j2}\beta_2 + \cdots + a_{jn}\beta_n &= -a_{j0}. \end{aligned}$$

Let us consider α and β as column vectors and suppose that α and β are linearly independent over \mathbb{F}_p .

Then there exist a pair (i, j) such that $\begin{vmatrix} \alpha_i & \alpha_j \\ \beta_i & \beta_j \end{vmatrix} \neq 0$. Applying the necessary permutation if needed,

we may assume that $\begin{vmatrix} \alpha_2 & \alpha_3 \\ \beta_2 & \beta_3 \end{vmatrix} \neq 0$. Then consider the invertible $(n-1) \times (n-1)$ matrix B

$$B := \begin{pmatrix} \alpha_2 & \alpha_3 & \cdots & \cdots & \alpha_n \\ \beta_2 & \beta_3 & \cdots & \cdots & \beta_n \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}$$

Let a_j denotes $a_j = (a_{22} \cdots a_{2n})^T$ for $2 \leq j \leq n$. Let $a_2 := B^{-1}(-1 \ 0 \ \cdots \ 0)^T$, so that $a_2 \cdot \alpha^T = -1$, $a_2 \cdot \beta^T = 0$. Let $a_3 := B^{-1}(1 \ e_1^T)^T$ so that $a_3 \cdot \alpha^T = 1$, $a_3 \cdot \beta^T = 1$ and let $a_j := B^{-1}(0 \ e_{j-2}^T)^T$ so that $a_j \cdot \alpha^T = 0$, $a_j \cdot \beta^T = 0$ for $3 < j < n$ where e_i 's denote canonical basis vectors for \mathbb{Z}_p^{n-2} . Let also $a_{20} = 1$, $a_{30} = -1$ and $a_{j0} = 0$. Now, if we define $A = (a_2 \cdots a_n)^T$ then by construction of a_i 's we have

$$A \cdot \alpha = \begin{pmatrix} a_2 \cdot \alpha^T \\ a_3 \cdot \alpha^T \\ \vdots \\ a_n \cdot \alpha^T \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \text{ and } A \cdot \beta = \begin{pmatrix} a_2 \cdot \beta^T \\ a_3 \cdot \beta^T \\ \vdots \\ a_n \cdot \beta^T \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}. \text{ Hence we get } a_2 \cdot \alpha^T + a_{20} =$$

$-1 + 1 = 0$ and $a_2 \cdot \beta^T + a_{20} = 0 + 1 = 1$ as needed. Also $a_3 \cdot \alpha^T + a_{30} = 1 - 1 = 0$ and $a_3 \cdot \beta^T + a_{30} = 1 - 1 = 0$ as needed. Also $a_j \cdot \alpha^T + a_{j0} = 0 + 0 = 0$ and $a_j \cdot \beta^T + a_{j0} = 0 + 0 = 0$ as needed. It remains to show that the set $\{a_2, a_3, \dots, a_n\}$ is linearly independent.

Now since B is invertible the set $\{a_2, a_3, \dots, a_n\}$ is linearly independent iff the set $\{Ba_2, Ba_3, \dots, Ba_n\}$ is linearly independent. Let for some $\gamma_i \in \mathbb{F}_p$, $2 \leq i \leq n$ we have

$$\gamma_2 \begin{pmatrix} -1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \gamma_3 \begin{pmatrix} 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \gamma_4 \begin{pmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} + \cdots + \gamma_n \begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = 0.$$

Then it can be easily seen that $\gamma_i = 0$ and hence $\{a_2, a_3, \dots, a_n\}$ is a linearly independent set. It follows that A is invertible and the translation we have constructed

$$\begin{pmatrix} a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots \\ a_{n2} & \dots & a_{nn} \end{pmatrix} \begin{pmatrix} x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

satisfies the conditions we needed.

Example: Let $\alpha = (2, 0, 0, 0)$ and $\beta = (2, 3, 0, 1)$ in \mathbb{Z}_5^4 . Then

$$B = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 2 & 3 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \Rightarrow B^{-1} = \begin{pmatrix} 3 & 0 & 0 & 0 \\ 3 & 2 & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \pmod{5}.$$

$$a_2 = B^{-1} \begin{pmatrix} -1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 0 \\ 0 \end{pmatrix}, a_3 = B^{-1} \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ 0 \\ 0 \end{pmatrix}, a_4 = B^{-1} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, a_5 =$$

$$B^{-1} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 3 \\ 0 \\ 1 \end{pmatrix}. \text{ Then the transformation is}$$

$$\begin{pmatrix} 2 & 2 & 0 & 0 \\ 3 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 3 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}$$

and $\bar{f}(x_1, x_2, x_3, x_4, x_5) = f(x_1, 1 + 2x_2 + 2x_3, -1 + 3x_2, x_4, 3x_3 + x_5)$ and

$$\begin{aligned} \bar{f}(x_1, 2, 0, 0, 0) &= f(x_1, 0, 0, 0, 0) \\ \bar{f}(x_1, 2, 3, 0, 1) &= f(x_1, 1, 0, 0, 0). \end{aligned}$$

Now suppose that α and β are linearly dependent over \mathbb{F}_p . Again applying the necessary permutation if needed, we may assume that $0 \neq \alpha_2$ and $\alpha_2 \neq \beta_2$. Let $a_{20} = -\alpha_2/(\beta_2 - \alpha_2)$, $a_{22} = 1/(\beta_2 - \alpha_2)$ and $a_{2j} = 0$ for $3 \leq j \leq n$. Then we have $a_2 \cdot \alpha^T + a_{20} = \frac{\alpha_2}{\beta_2 - \alpha_2} - \frac{\alpha_2}{\beta_2 - \alpha_2} = 0$ and $a_2 \cdot \beta^T + a_{20} = \frac{\beta_2}{\beta_2 - \alpha_2} - \frac{\alpha_2}{\beta_2 - \alpha_2} = 1$ as needed.

Now consider the $1 \times (n-1)$ matrix $B = (\alpha_2 \dots \alpha_n)$. Since $\alpha \neq 0$ $\dim(\text{Ker}(B)) = n-2$. Let v_2, \dots, v_n be a basis for $\text{Ker}(B)$. Let for some $\gamma_i \in \mathbb{F}_p$, $2 \leq i \leq n$ we have

$$\gamma_2 a_2 + \gamma_3 v_3 + \gamma_4 v_4 + \dots + \gamma_n v_n = 0.$$

Then applying B from left hand side we have $\gamma_2 B a_2 = \gamma_2 \alpha_2 / (\beta_2 - \alpha_2) = 0 \Rightarrow \gamma_2 = 0$. It follows that $\gamma_i = 0$ and $\{a_2, v_2, \dots, v_n\}$ is linearly independent. Then $A := (a_2 v_2 \dots v_n)^T$ is invertible. It can be

readily verified that the translation we have constructed

$$A \begin{pmatrix} x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} \frac{-\alpha_2}{\beta_2 - \alpha_2} \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

satisfies the remaining conditions needed.

Example: Let $\alpha = (1, 1, 0, 0)$ and $\beta = (2, 2, 0, 0)$ in \mathbb{Z}_5^4 . Then $a_{20} = -\alpha_2/(\beta_2 - \alpha_2) = 4$ and $a_{22} = 1/(\beta_2 - \alpha_2) = 1 \pmod{5}$. $B = (1\ 1\ 0\ 0)$

$$v_3 = \begin{pmatrix} 3 \\ 2 \\ 0 \\ 0 \end{pmatrix}, v_4 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, v_5 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

and then the transformation is

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 3 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} + \begin{pmatrix} 4 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

where $\bar{f}(x_1, x_2, x_3, x_4, x_5) = f(x_1, 4 + x_2, 3x_2 + 2x_3, x_4, x_5)$ and

$$\begin{aligned} \bar{f}(x_1, 1, 1, 0, 0) &= f(x_1, 0, 0, 0, 0) \\ \bar{f}(x_1, 2, 2, 0, 0) &= f(x_1, 1, 0, 0, 0). \end{aligned}$$

□

2.1 A comparison with the binomial distribution.

Let Y be a random variable from a binomial distribution $B(n, p)$ with n trials and probability p . So $0 \leq Y \leq n$, $\text{Prob}[Y = k] = \binom{n}{k} p^k (1-p)^{n-k}$, $E[Y] = np$ and $\text{Var}[Y] = np(1-p)$. We noticed that the mean and variance of X in Theorem 2 is the same as the mean and variance of the binomial distribution $B(n, p)$ with $n = q$ trials and probability $p = 1/q$. In Table 1 below we compare the two distributions for

$$\begin{aligned} f &= x^2 + (a_1y + a_2)x + (a_3y^2 + a_4y + a_5) \text{ and} \\ g &= x^2 + (b_1y + b_2)x + (b_3y^2 + b_4y + b_5) \end{aligned}$$

in $\mathbb{F}_q[x, y]$ with $q = 7$. Note that there are 7^{10} pairs for f, g . In Table 1 F_k is the number of pairs for which $\text{gcd}(f(x, \alpha), g(x, \alpha)) \neq 1$ for exactly k values for $\alpha \in \mathbb{F}_7$. We computed F_k by computing this gcd for all distinct pairs using Maple. The values for B_k come from $B(7, 1/7)$. They are given by $B_k = 7^{10} \text{Prob}[Y = k]$.

k	0	1	2	3	4	5	6	7
F_k	96606636	110666892	56053746	17287200	1728720	0	0	132055
B_k	96018048	112021056	56010528	15558480	2593080	259308	14406	343

Table 1: Data for quadratic (f, g) in $\mathbb{F}_7[x, y]$

The two zeros F_5 and F_6 can be explained as follows. Let $R(y)$ be the Sylvester resultant of f and g . Then applying Lemma 1 we have $R(\alpha) = 0 \iff \gcd(f(x, \alpha), g(x, \alpha)) \neq 1$ for $\alpha \in \mathbb{F}_q$. For our quadratic polynomials f and g , Lemma 1(ii) implies $\deg R \leq \deg f \deg g = 4$. Hence $R(y)$ can have at most 4 distinct roots unless f and g are not coprime in $\mathbb{F}_7[x, y]$ in which case $R(y) = 0$ and it has 7 roots. Therefore $F_5 = 0$, $F_6 = 0$ and $F_7 = 132055$ is the number pairs f, g which are not coprime in $\mathbb{F}_7[x, y]$.

References

- [1] Arthur Benjamin and Curtis Bennett. The Probability of Relatively Prime Polynomials. *Mathematics Magazine* **80** (3), 197–202, 2007.
- [2] Erwin Berlekamp. *Algebraic Coding Theory*, McGraw-Hill, 1968.
- [3] David Cox, John Little and Donal O’Shea. *Ideals, Varieties and Algorithms*. Springer-Verlag, 3rd ed., 2007.
- [4] J. de Kleine, M. B. Monagan, A. D. Wittkopf. Algorithms for the Non-monic case of the Sparse Modular GCD Algorithm. *Proc. ISSAC ’05*, ACM Press, (2005), 124–131.
- [5] Arnold Knopfmacher and John Knopfmacher. Counting irreducible factors of polynomials over finite fields. *Discrete Mathematics* **112** (1993) 103–118.
- [6] Michael Monagan and Baris Tuncer. Using Sparse Interpolation to Solve Multivariate Diophantine Equations. *Communications in Computer Algebra* 49:3, Issue 193, pp. 94–97, September 2015.
- [7] Sequence <http://oeis.org/A006579> in *The On-Line Encyclopedia of Integer Sequences*, published electronically at <http://oeis.org>, 2010.
- [8] Gary Mullen and Daniel Panario. 11.2.3.4 Greatest common divisors of polynomials. *Handbook of Finite Fields*. CRC Press, 2013.
- [9] Wolfgang Schmidt. *Equations over Finite Fields: An Elementary Approach*. Springer-Verlag LNCS **536** (1976) Ch 4 pp. 157–159.
- [10] Jack Schwartz, Fast probabilistic algorithms for verification of polynomial identities. *J. ACM* **27** 701–717, 1980. See also https://en.wikipedia.org/wiki/Schwartz%E2%80%93Zippel_lemma.
- [11] Richard. Zippel. Probabilistic Algorithms for Sparse Polynomials, *Proc. EUROSAM ’79*, Springer-Verlag LNCS, **2**, 216–226, 1979.

Appendix A

Below is Magma code for quadratic polynomials over \mathbb{F}_4 . For each pair of quadratic polynomials $F, G \in \mathbb{F}_4[x, y]$ we compute $X = |\{\alpha \in \mathbb{F}_4 : \gcd(F(x, \alpha), G(x, \alpha)) \neq 1\}|$. The code counts A_k the number of pairs (F, G) with $k = X$ and computes $E[X]$ and $\text{Var}[X]$.

```

q := 4;
Fq<z> := FiniteField(q);
P<x,y> := PolynomialRing(Fq,2);
N := 0; // counter
M := 0; // mean
V := 0; // variance
A := AssociativeArray(); // frequencies

for X in [0..q] do A[X] := 0; end for;

for a in Fq do for b in Fq do
for c in Fq do for d in Fq do for e in Fq do
for r in Fq do for s in Fq do
for t in Fq do for u in Fq do for v in Fq do

if not ( [a,b,c,d,e] gt [r,s,t,u,v] ) then
  X := 0;
  for y in Fq do
    F := x^2+(a*y+b)*x+(c*y^2+d*y+e);
    G := x^2+(r*y+s)*x+(t*y^2+u*y+v);
    if Gcd(F,G) ne 1 then X := X+1; end if;
  end for;
  if [a,b,c,d,e] eq [r,s,t,u,v] then
    N := N+1; A[X] := A[X]+1;
    M := M+X; V := V+(X-1)*(X-1);
  else
    N := N+2; A[X] := A[X]+2;
    M := M+2*X; V := V+2*(X-1)*(X-1);
  end if;
end if;

end for; end for; end for; end for; end for;
end for; end for; end for; end for; end for;

"field size", q;
"N", N, q^10;
"frequencies", A[0],A[1],A[2],A[3],A[4];
"mean", 1.0*M/q^10;
"variance", 1.0*V/q^10;

```