Research Problems

Edited by P. Borwein, I. Daubechies, and V. Totik
(*Please report progress on the problems to Research_Problems@cecm.sfu.ca*)

Constr. Approx. (1996) 12: 555–558

Bivariate Segment Approximation and Free Knot Splines: Research Problems 96-4

G. Nürnberger

In this note we state some problems on approximation by univariate splines with free knots, bivariate segment approximation, and tensor product splines with variable knot lines.

There is a vast literature on approximation and interpolation by univariate splines with fixed knots (see, e.g., the books by de Boor [1], Braess [2], DeVore and Lorentz [4], Powell [20], Schumaker [21], Nürnberger [13], and the book by Chui [3] on multivariate splines). On the other hand, numerical examples show that, in general, the error is much smaller if variable knots are used for the approximation of functions instead of fixed knots. This is true for univariate splines as well as for bivariate splines. But approximation by splines with free knots leads to rather difficult nonlinear problems.

We first consider best approximation by univariate splines with free knots. We denote by $S_{m,k}$ the (nonconvex) set of *splines* of degree m with k *free knots*. Given a function $f \in C[a, b]$, we call $s_f \in S_{m,k}$ a *best approximation* of f if $\|f - s\| = \min_{s \in S_{m,k}} \|f - s\|$, where $\|h\| = \max_{t \in [a,b]} |h(t)|$ denotes the *maximum norm* of $h \in C[a, b]$. This problem was first investigated by Johnson [5] in 1960. Since then, many results were proved on this topic by using advanced methods (see, e.g., the books [2], [4], [13], [21] and the recent results in [6], [10], [11], [12], [14], [15], [16], [17], [22]). Nevertheless, there is no complete solution of the following problem.

Problem 1. *Give characterizations of best approximations and unique best approximations from* $S_{m,k}$.

It is well known that, in contrast to the fixed knot case, best approximations from $S_{m,k}$ cannot be characterized by alternation properties of the error function alone. But at present, it seems not to be known what the additional conditions could be which completely characterize best approximations.

The deeper reason that nonlinear problems of this type are unsolved is that there is no complete theory of nonlinear optimization and no algorithm for computing global minima of arbitrary nonlinear optimization problems.

On the other hand, an algorithm was developed for computing good spline approximations with free knots (Nürnberger, Sommer, and Strauss [19], and Meinardus, Nürnberger, Sommer, and Strauss [8]). The spline approximations are computed in two steps.

Given a function $f \in C[a, b]$, first a segment approximation problem is solved. We denote by $PP_{m,k}$ the (nonconvex) set of *piecewise polynomials* of degree m with k *free*

knots. In the first step, a best approximation of f from $PP_{m,k}$ is computed such that the errors on all knot intervals are the same. Therefore, the corresponding knots, denoted by $x_1, \ldots, x_k$ (which, in general, are nonuniform), reflect the critical parts of the function f. The algorithm converges by starting with an arbitrary set of knots.

Then in the second step, by applying a Remez-type algorithm (Nürnberger and Sommer [18]), a best approximation of f from the space $S_m(x_1, \ldots, x_k)$ of *splines* of degree m with k *fixed knots* is computed. Numerical examples show that in this way a good approximation from $S_{m,k}$ is obtained (see [8] and [13]). The resulting error can be compared with the error for $PP_{m,k}$ which is a lower bound.

In the following, we consider similar problems in the bivariate case. First, we formulate a general spline approximation problem for variable knot lines.

Let a rectangle $T = [a, b] \times [c, d]$ be subdivided by knot lines $x = x_i$ and $y = y_i$, $i = 1, \ldots, k$, into $(k+1)^2$ subrectangles. Such a *partition* is called *of type* I. Moreover, let an approximation or interpolation method be given which yields for each function $f \in C(T)$ an approximation $A(f)$ from the tensor product spline space $S_m(x_1, \ldots, x_k) \otimes S_m(y_1, \ldots, y_k)$ (see, e.g., [1] and [13]). We consider the case when the knot lines are variable.

Problem 2. *Determine a partition of type* I *for which the error* $\|f - A(f)\|_T$ *is* (*nearly*) *minimal*.

As in the univariate case, it is natural to get a good partition as in Problem 2 by solving segment approximation problems for $\Pi_n \otimes \Pi_n$ (where Π_n denotes the space of univariate *polynomials* of degree n) of the following type.

We subdivide the rectangle T by k *horizontal* lines into $k+1$ strips. Then we subdivide each strip by k vertical line segments into $k+1$ subrectangles. The partitions of each strip are different, in general. Such a *partition* of T is called *of type* II. If we subdivide T by k *vertical* lines into $k+1$ strips, and subdivide each strip by k horizontal line segments into $k+1$ subrectangles, then the *partition* is called *of type* III.

We associate to each subrectangle T_j of a partition of type I, II, or III a real number $d(T_j)$ which may be the error $\|f - B(f)\|_{T_j}$ of some approximation or interpolation method, where $B(f)$ is from the tensor product space $\Pi_n \otimes \Pi_n$. The real number may also be a lower or upper bound of this error. We consider the case when the line segments of the partition are variable.

Problem 3. *Describe and determine the partitions of type* I, II, *or* III *for which* $\max_j d(T_j)$ *is minimal*.

There may be relations between optimal partitions of type I, II, and III.

Problem 4. *For which functionals* d *do the horizontal lines of an optimal partition of type* II *combined with the vertical lines of an optimal partition of type* III *yield an optimal partition of type* I (*in the sense of Problem* 3)?

In a recent paper, Meinardus, Nürnberger, and Walz [9] developed an algorithm for computing optimal partitions of type II and III for the best approximation error and

related functionals d (Problem 3). They also showed that for some of these functionals, Problem 4 has a positive answer. Their numerical examples show that tensor product polynomial interpolation at uniform points on the subrectangles is a suitable method for segment approximation (although this method does not quite fit into the general setting of the convergence results). An optimal partition of type I in the sense of Problem 3 yields a good partition for tensor product spline interpolation in the sense of Problem 2.

Although, first results on the above problems are known, several problems for various approximation methods with different function classes are unsolved at present. Efficient methods for solving these problems would be important for applications in practice.

References

1. C. DE BOOR (1978): A Practical Guide to Splines. New York: Springer-Verlag.
2. D. BRAESS (1986): Nonlinear Approximation Theory. Berlin: Springer-Verlag.
3. C. K. CHUI (1988): Multivariate Splines. Philadelphia: CBMS-SIAM.
4. R. A. DEVORE, G. G. LORENTZ (1993): Constructive Approximation. New York: Springer-Verlag.
5. R. S. JOHNSON (1960): *On monosplines of least deviation*. Trans. Amer. Math. Soc., **96**:458–477.
6. H. KAWASAKI (1994): *A second-order property of spline functions with one free knot*. J. Approx. Theory, **78**:293–297.
7. G. MEINARDUS (1967): Approximation of Functions: Theory and Numerical Methods. Berlin: Springer-Verlag.
8. G. MEINARDUS, G. NÜRNBERGER, M. SOMMER, H. STRAUSS (1989): *Algorithms for piecewise polynomials and splines with free knots*. Math. Comput., **53**:235–247.
9. G. MEINARDUS, G. NÜRNBERGER, G. WALZ (to appear): *Bivariate segment approximation and splines*. Advances in Computational Mathematics.
10. B. MULANSKY (1992): *Chebyshev approximation by spline functions with free knots*. IMA J. Numer. Anal., **12**:95–105.
11. B. MULANSKY (1992): *Necessary conditions for local best Chebyshev approximations by splines with free knots*. In: Numerical Methods of Approximation Theory (D. Braess, L. L. Schumaker, eds.). ISNM, 105. Basel: Birkäuser, pp. 195–206.
12. G. NÜRNBERGER (1987): *Strongly unique spline approximation with free knots*. Constr. Approx., **3**:31–42.
13. G. NÜRNBERGER (1989): Approximation by Spline Functions. Berlin: Springer-Verlag.
14. G. NÜRNBERGER (1992): *The metric projection for free knot splines*. J. Approx. Theory, **71**:145–153.
15. G. NÜRNBERGER (1994a): *Strong unicity in nonlinear approximation and free knot splines*. Constr. Approx., **10**:285–299.
16. G. NÜRNBERGER (1994b): *Approximation by univariate and bivariate splines*. In: Second International Colloquium on Numerical Analysis (D. Bainov, V. Covachev, eds.). Utrecht: VSP, pp. 143–153.
17. G. NÜRNBERGER, L. L. SCHUMAKER, M. SOMMER, H. STRAUSS (1989): *Uniform approximation by generalized splines with free knots*. J. Approx. Theory, **59**:150–169.
18. G. NÜRNBERGER, M. SOMMER (1983): *A Remez type algorithm for spline functions*. Numer. Math., **41**:117–146.
19. G. NÜRNBERGER, M. SOMMER, H. STRAUSS (1986): *An algorithm for segment approximation*. Numer. Math., **48**:463–477.
20. M. J. D. POWELL (1981): Approximation Theory and Methods. Cambridge, UK: Cambridge University Press.
21. L. L. SCHUMAKER (1981): Spline Functions: Basic Theory. New York: Wiley-Interscience.

22. M. SOMMER (1992): *Segment approximation using linear functionals*. In: Numerical Methods of Approximation Theory (D. Braess, L. L. Schumaker, eds.). ISNM, 105. Basel: Birkäuser, pp. 331–346.

G. Nürnberger
Fakultät für Mathematik und Informatik
Universität Mannheim
D-68131 Mannheim
Germany