

**SOME OLD PROBLEMS
ON POLYNOMIALS WITH
INTEGER COEFFICIENTS**

PETER BORWEIN

Simon Fraser University Centre for
Constructive and Experimental
Mathematics

<http://www.cecm.sfu.ca/~pborwein>

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

- Some old chestnuts. All at least 35 years old.
- All involve Chebyshev type problems for polynomials with integer coefficients.
- All very hard.
- All have a highly non-trivial computational component.
- All have accessible partials?
- All very interesting.

A. The Integer Chebyshev Problem of Hilbert and Fekete.

1. Problem. Find

$$C_N[\alpha, \beta] := \left(\min_{a_i \in \mathbb{Z}, a_N \neq 0} \|a_0 + a_1x + \dots + a_Nx^N\|_{[\alpha, \beta]} \right)^{\frac{1}{N}}.$$

We will restrict to $\beta - \alpha \leq 4$.

2. One can show that

$$C[\alpha, \beta] := \lim_{N \rightarrow \infty} C_n[\alpha, \beta]$$

exists. This is the integer Chebyshev constant for the interval or the integer transfinite diameter.

3. We (T. Erdélyi and P.B.) show

$$\frac{1}{2.3768 - \epsilon} \leq C[0, 1] \leq \frac{1}{2.360}.$$

Lemma. *Suppose*

$$q_m(x) = a_m x^m + \dots + a_0, \quad a_m \in \mathbb{Z}$$

has all its roots in $(0, 1)$. (That is: $q_m \in TR(0, 1)$). Then, provided $(q_m, p_n) = 1$

$$\|p_n\|_{[0,1]}^{1/n} \geq \frac{1}{a_m^{1/m}}.$$

- It is conjectured (Chudnovskys, Montgomery) that the lemma gives the right bound in 3. (This is likely false.)

B. The Schur, Siegel, Smyth Trace Problem.

1. Conjecture. Suppose

$$p_n(z) = a_n z^n + \dots + a_0, a_i \in \mathbb{Z}$$

has all real, positive roots and is irreducible.
Then

$$a_{n-1} \geq (2 - \epsilon)n.$$

2. Partial. Except for finitely many (explicit) exceptions

$$a_{n-1} \geq e^{1/2}n \quad \text{Schur (1918)}$$

$$a_{n-1} \geq (1.733\dots)n \quad \text{Siegel (1943)}$$

$$a_{n-1} \geq (1.771\dots)n \quad \text{Smyth (1983).}$$

3. The Relationship to the Small Interval Problem.

Lemma. *If*

$$C[0, 1/m] \leq 1/(m + \delta)$$

then, for totally positive polynomials

$$a_{n-1} \geq \delta n$$

(with finitely many explicit exceptions).

Corollary. $\delta > 1.744$

Proof. By example on $C[0, 1/100]$.

C. Prouhet-Tarry-Escott Problem.

1. Conjecture. For any N there exists $p \in Z[x]$ (a polynomial with integer coefficients) so that

$$p(x) = (x - 1)^N q(x) = \sum a_k x^k$$

and

$$S(p) := \sum |a_k| = 2N.$$

Almost equivalently (though not quite obviously) this polynomial must have coefficients $\{0, -1, +1\}$ and so

$$\|p\|_{L^2\{|z|=1\}} = \sqrt{2N}.$$

2. The Basis for the Conjecture.

$$x^{\alpha_1} + \dots + x^{\alpha_N} - x^{\beta_1} - \dots - x^{\beta_N} = 0((x-1)^N).$$

For $N = 2, \dots, 10$ with

$$[\alpha_1, \dots, \alpha_N] \quad \text{and} \quad [\beta_1, \dots, \beta_N]$$

- $[0, 3] = [1, 2]$
- $[1, 2, 6] = [0, 4, 5]$
- $[0, 4, 7, 11] = [1, 2, 9, 10]$
- $[1, 2, 10, 14, 18] = [0, 4, 8, 16, 17]$
- $[0, 4, 9, 17, 22, 26] = [1, 2, 12, 14, 24, 25]$
- $[0, 18, 27, 58, 64, 89, 101]$
 $= [1, 13, 38, 44, 75, 84, 102]$

- $[0, 4, 9, 23, 27, 41, 46, 50]$

$= [1, 2, 11, 20, 30, 39, 48, 49]$

- $[0, 24, 30, 83, 86, 133, 157, 181, 197]$

$= [1, 17, 41, 65, 112, 115, 168, 174, 198]$

- $[0, 3083, 3301, 11893, 23314, 24186, 35607,$

$44199, 44417, 47500] =$

$[12, 2865, 3519, 11869, 23738, 23762, 35631,$

$43981, 44635, 47488]$

- The size 10 example illustrates the problems inherent with searching for a solution.

3. Partial History.

- Euler
- Prouhet (1851)
- Tarry (1910) - Small Examples
- Escott (1910) - Small Examples
- Letac (1941) - Size 9 and 10
- Gloden (1946) - Size 9 and 10
- Smyth (Math Comp. 1991) - Size 10 generalized.

4. Diophantine Form

Find distinct integers

$$[\alpha_1, \dots, \alpha_N] \text{ and } [\beta_1, \dots, \beta_N]$$

so that

$$\alpha_1 + \dots + \alpha_N = \beta_1 + \dots + \beta_n$$

$$\alpha_1^2 + \dots + \alpha_N^2 = \beta_1^2 + \dots + \beta_n^2$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$\alpha_1^{N-1} + \dots + \alpha_N^{N-1} = \beta_1^{N-1} + \dots + \beta_N^{N-1}$$

- The problem is completely open for $N \geq 11$.

D. The Weak Prouhet-Tarry-Escott Problem.

1. Problem. For fixed N find $p \in Z[x]$

$$p(x) = (x - 1)^N q(x) = \sum a_k x^k$$

that minimizes

$$S(p) = \sum |a_i|$$

or

$$S^2(p) = (\sum |a_i|^2)^{1/2}$$

2. Solving $S(p) = |S^2(p)|^2 = 2N$ is the Prouhet-Tarry-Escott-Problem and is the big prize.

3. Showing that there exist

$$\{p_N\} = \{(x-1)^N q(x)\}$$

so that

$$S(p_N) = o(N \log N)$$

is also a big prize.

- This shows that the “Easier Waring Problem” is easier than the “Waring Problem” (At the moment.)
- That is: it requires essentially fewer powers to write every integer as sums and differences of *Nth* powers than just as sums of *Nth* powers. (Fuchs and Wright, Quart. J. Math. 1936).

4. It is known that

$$S((x-1)^N q(x)) \leq \frac{N^2}{2}$$

is possible.

Any improvement would be a major step.

5. If we demand that p has a zero of order N but not $N+1$ at 1 then

$$S(p) = O((\log N)N^2)$$

is possible (Hua).

Any improvement would be interesting.

E. Problem of Erdős and Szekeres (1958).

1. Problem. Minimize over $\{\alpha_1, \dots, \alpha_N\}$

$$S \left(\prod_{k=1}^N (1 - x^{\alpha_i}) \right)$$

Call this minimum S_N^π .

2. Conjecture. $S_N^\pi \gg N^k$ for any k .

3. From the P-T-E problem

$$S_N^\pi \geq 2N$$

4. Examples.

N	$\ f\ _1$	$\{\alpha_1, \dots, \alpha_N\}$
1	2	$\{1\}$
2	4	$\{1, 2\}$
3	6	$\{1, 2, 3\}$
4	8	$\{1, 2, 3, 4\}$
5	10	$\{1, 2, 3, 5, 7\}$
6	12	$\{1, 1, 2, 3, 4, 5\}$
7	16	$\{1, 2, 3, 4, 5, 7, 11\}$
8	16	$\{1, 2, 3, 5, 7, 8, 11, 13\}$
9	20	$\{1, 2, 3, 4, 5, 7, 9, 11, 13\}$
10	24	$\{1, 2, 3, 4, 5, 7, 9, 11, 13, 17\}$
11	28	$\{1, 2, 3, 5, 7, 8, 9, 11, 13, 17, 19\}$
12	36	$\{1, \dots, 9, 11, 13, 17\}$
13	48	$\{1, \dots, 9, 11, 13, 17, 19\}$

5. Conjecture. Except for $N = 1, 2, 3, 4, 5, 6$ and 8

$$S_N^\pi \geq 2N + 2.$$

- Maltby solves this for $N=7, 9$ and 10.

6. Partials.

$$S_N^\pi \ll N^{0(N^{1/2})} \quad (\text{Atkinson, Dobrowolski})$$

$$S_N^\pi \ll N^{0(\log N N^{1/3})} \quad (\text{Odlyzko})$$

(could equally well use $\| \cdot \|_{L^2(D)}$.)

F. Lehmer's Conjecture.

Mahler's Measure: if

$$p(z) = \prod_{i=1}^n (z - \alpha_i)$$

then

$$M(p) = \prod_{i=1}^n \max\{1, |\alpha_i|\}$$

or equivalently

$$M(p) := \exp \left\{ \int_0^1 \log |p(e^{2\pi it})| dt \right\}$$

Conjecture. Suppose p is a monic polynomial with integer coefficients. Then either $M(p) = 1$ or $M(p) > 1.17\dots$

- This generalizes Kronecker's theorem which can be stated as: $M(p) = 1$ implies that p is cyclotomic.

- The minimal Mahler measure for a non-cyclotomic p is speculated to be

$$p := x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1$$

for which $M(p) = 1.17628081825991750\dots$

- This is also speculated to be the smallest Salem number.

Question. Do there polynomials with coefficients $\{0, -1, +1\}$ with roots of arbitrarily high multiplicity inside the unit disk.

A negative answer solves the conjecture.

G. Littlewood's Conjecture.

Conjecture (resolved 1981). Suppose

$$p(z) := \sum_{n=0}^N c_n z^{K_n} \quad \|c_i\| \geq 1.$$

Then the L_1 norm of p on the boundary of the unit disc is $\gg \log(N)$.

H. A Problem of Erdős.

Conjecture, 1957. Suppose

$$p(z) := \sum_{n=0}^N c_n z^n \quad c_i \pm 1.$$

Then the supremum norm of p on the boundary of the unit disc is $> (1 + \epsilon)\sqrt{N}$.

I. Littlewood's Other Conjecture.

Conjecture (1966). There is some

$$p(z) := \sum_{n=0}^N c_n z^n \quad c_i \pm 1$$

so that for all z on the boundary of the unit disc

$$C_1 < \frac{|p(z)|}{\sqrt{n}} < C_2.$$

- Littlewood, in part, based his conjecture on computations of all such polynomials up to degree twenty.
- Odlyzko has now done 200 MIPS years of computing on this problem

SO MUCH FOR MOTIVATION

**ON THE NUMBER OF
ZEROS OF $\{0, +1, -1\}$
POLYNOMIALS AND RELATED
CHEBYSHEV PROBLEMS.**

P. BORWEIN, T. ERDÉLYI AND G. KÓS

- We consider the problem of minimizing the uniform norm on $[0, 1]$ over polynomials p

$$p(x) = \sum_{j=m}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}$$

with fixed $|a_m| \neq 0$.

- This is equivalent to the question of how many zeros such a polynomial can have at 1.

- Particular cases include:

Polynomials with coefficients in the set $\{-1, 0, 1\}$.

Polynomials with coefficients in the set $\{0, 1\}$ on the interval $[-1, 0]$.

$$\mathcal{P}_n := \left\{ \sum_{i=0}^n a_i x^i : a_i \in \mathbb{R} \right\}$$

$$\mathcal{Z}_n := \left\{ \sum_{i=0}^n a_i x^i : a_i \in \mathbb{Z} \right\}$$

$$\mathcal{F}_n := \left\{ \sum_{i=0}^n a_i x^i : a_i \in \{-1, 0, 1\} \right\}$$

$$\mathcal{A}_n := \left\{ \sum_{i=0}^n a_i x^i : a_i \in \{0, 1\} \right\}$$

So obviously

$$\mathcal{A}_n \subset \mathcal{F}_n \subset \mathcal{Z}_n \subset \mathcal{P}_n.$$

2. NUMBER OF ZEROS AT 1

Theorem 2.1. *There is an absolute constant $c > 0$ such that every polynomial p of the form*

$$p(x) = \sum_{j=0}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}$$

has at most

$$c (n(1 - \log |a_0|))^{1/2}$$

zeros at 1.

- Applying Theorem 2.1 with $q(x) := x^{-n}p(x^{-1})$ gives the following:

Theorem 2.2. *There is an absolute constant $c > 0$ such that every polynomial p of the form*

$$p(x) = \sum_{j=0}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}$$

has at most

$$c (n(1 - \log |a_n|))^{1/2}$$

zeros at 1.

- This sharpens a sequence of old and not so old results of Littlewood, Schur, Turán, Erdős, Bombieri, Vaaler and others. (In the case of small height polynomials.)
- The result is sharp.

Theorem 2.3. *It $\exp(-3n) \leq |a_0| \leq 1$, then there exists a polynomial p of the form*

$$p(x) = \sum_{j=0}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}$$

such that p has a zero at 1 with multiplicity at least

$$\frac{1}{5}(n(1 - \log |a_0|))^{1/2} - 1.$$

- The next two theorems treat the case $a_0 = 1$. The proofs are attractive and we will work through them. (As time allows.)

Theorem 2.4. *Every polynomial p of the form*

$$p(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| = 1, \quad |a_j| \leq 1$$

has at most $5\sqrt{n}$ zeros at 1 and at most $c\sqrt{n}$ zeros in $[-1, 1]$.

Theorem 2.5. *For every $n \in \mathbb{N}$, there exists*

$$p_n(x) = \sum_{j=0}^{n^2} a_j x^j$$

such that $a_0 = 1$; a_1, a_2, \dots, a_{n^2} are real numbers of modulus less than 1; and p_n has a zero at 1 with multiplicity at least $n - 1$.

- Theorem 2.5 immediately implies

Corollary 2.6. *For every $n \in \mathbb{N}$, there exists a polynomial*

$$p_n(x) = \sum_{j=0}^n a_j x^j, \quad a_n = 1,$$

a_0, a_1, \dots, a_n are real numbers of modulus less than 1, and p_n has a zero at 1 with multiplicity at least $\lfloor \sqrt{n} \rfloor - 1$.

- The next related result is well known:

Theorem 2.7. *There is an absolute constant $c > 0$ so that for every $n \in \mathbb{N}$ there is a $p \in \mathcal{F}_n$ having at least $c\sqrt{n/\log(n+1)}$ zeros at 1.*

- Theorems 2.4 and 2.7 show that the right upper bound for the number of zeros a polynomial $p \in \mathcal{F}_n$ can have at 1 is somewhere between $c_1\sqrt{n/\log(n+1)}$ and $c_2\sqrt{n}$ with absolute constants $c_1 > 0$ and $c_2 > 0$.
- This gap looks quite hard to close. This is an old problem on which there has been no progress in 20 years.

- There is a simple observation about the maximal number of zeros a polynomial $p \in \mathcal{A}_n$ can have.

Theorem 2.8. *There is an absolute constant $c > 0$ such that every $p \in \mathcal{A}_n$ has at most $c \log n$ zeros at -1 .*

- There is a less simple observation about the maximal number of zeros at 1 of a polynomial with coefficients $\{+1, -1\}$.

There are between $c_1 \log n$ and $c_2(\log n)^2$ such zeros and is open as to what is correct (Boyd 95).

Remark to Theorem 2.8. Let R_n be defined by

$$R_n(x) := \prod_{i=1}^n (1 + x^{a_i}),$$

where $a_1 := 1$ and a_{i+1} is the smallest odd integer that is greater than $\sum_{k=1}^i a_k$.

- It is tempting to speculate that R_n is the lowest degree polynomial with coefficients $\{0, 1\}$ and a zero of order n at -1 .
- This is true for $n := 1, 2, 3, 4, 5$ but fails for $n := 6$ and hence for all larger n .

3. RESTRICTED CHEBYSHEV PROBLEM

Theorem 3.1. *There are absolute constants so that*

$$\begin{aligned} & \exp \left(-c_1 n (1 - \log |a_m|) \right)^{1/2} \\ & \leq \inf_p \|p\|_{[0,1]} \\ & \leq \exp \left(-c_2 n (1 - \log |a_m|) \right)^{1/2} , \end{aligned}$$

where the *inf* is taken over $0 \neq p$ of the form

$$p(x) = \sum_{j=m}^n a_j x^j, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}$$

with $|a_m| \geq \exp \left(\frac{1}{2} (1 - n) \right)$.

- This specializes to

Theorem 3.2. *There are absolute constants $c_1 > 0$ and $c_2 > 0$ such that*

$$\exp(-c_1\sqrt{n}) \leq \inf_p \|p\|_{[0,1]} \leq \exp(-c_2\sqrt{n}) ,$$

for polynomials of the form

$$p(x) = \sum_{j=m}^n a_j x^j , \quad |a_j| \leq 1 , \quad a_n = 1 .$$

- For the class \mathcal{F}_n we have

Theorem 3.3. *There are absolute constants $c_1 > 0$ and $c_2 > 0$ such that*

$$\begin{aligned} & \exp(-c_1 \sqrt{n}) \\ & \leq \inf_{0 \neq p \in \mathcal{F}_n} \|p\|_{[0,1]} \\ & \leq \exp(-c_2 \sqrt{n}). \end{aligned}$$

- The upper bound requires some new ideas.

- The approximation rate in Theorems 3.2 and 3.3 should be compared with

$$\min_{p(x) := x^n + \dots \in \mathcal{P}_n} \|p\|_{[0,1]}^{1/n} = \frac{2^{1/n}}{4},$$

and also with

$$\frac{1}{2.376\dots} < \min_{0 \neq p \in \mathcal{Z}_n} \|p\|_{[0,1]}^{1/n} < \frac{1 + o(1)}{2.3605}.$$

- The first equality above is attained by the normalized Chebyshev polynomial shifted linearly to $[0, 1]$ and is proved by a simple perturbation argument. The second inequality is much harder (the exact result is open).

- It is an interesting fact that the polynomials $0 \neq p \in \mathcal{Z}_n$ with the smallest uniform norm on $[0, 1]$ are very different from the usual Chebyshev polynomial of degree n .
- For example, they have at least 52% of their zeros at either 0 or 1. Relaxation techniques do not allow for their approximate computation.
- Likewise, polynomials $0 \neq p \in \mathcal{F}_n$ with small uniform norm on $[0, 1]$ are again quite different from polynomials $0 \neq p \in \mathcal{Z}_n$ with small uniform norm on $[0, 1]$.

- The story is roughly as follows. Polynomials $0 \neq p \in \mathcal{P}_n$ with leading coefficient 1 and with smallest possible uniform norm on $[0, 1]$ are characterized by equioscillation and are given by the Chebyshev polynomials explicitly.
- In contrast, finding polynomials from \mathcal{Z}_n with small uniform norm on $[0, 1]$ is closely related to finding irreducible polynomials with all their roots in $[0, 1]$.

- As we shall see the construction of small norm polynomials from \mathcal{F}_n is governed by how many zeros such a polynomial can have at 1.
- It is interesting to note that the polynomials $0 \neq p \in \mathcal{P}_n$ with leading coefficient 1 and with smallest uniform norm on $[0, 1]$ have coefficients that alternate in sign.
- This also appears to be true for the analogous polynomials from \mathcal{Z}_n (though this is only conjectural and probably quite hard to prove).

- This is quite different from the story for \mathcal{F}_n . For polynomials $p(-x)$ with $0 \neq p \in \mathcal{A}_n$ we get a very much larger smallest possible uniform norm on $[0, 1]$.

Theorem 3.4. *There are absolute constants $c_1 > 0$ and $c_2 > 0$ such that*

$$\begin{aligned} & \exp(-c_1 \log^2(n+1)) \\ & \leq \inf_{0 \neq p \in \mathcal{A}_n} \|p(-x)\|_{[0,1]} \\ & \leq \exp(-c_2 \log^2(n+1)) \end{aligned}$$

4. TOOLS

- In the general case the tools are:
- Denote by \mathcal{S} the collection of all analytic functions f on the open unit disk $D := \{z \in \mathbb{C} : |z| < 1\}$ that satisfy

$$|f(z)| \leq \frac{1}{1 - |z|}, \quad z \in D.$$

Theorem 4.1. *There are absolute constants $c_1 > 0$ and $c_2 > 0$ such that*

$$|f(0)|^{c_1/a} \leq \exp\left(\frac{c_2}{a}\right) \|f\|_{[1-a,1]}$$

for every $f \in \mathcal{S}$ and $a \in (0, 1]$.

Hadamard Three Circles Theorem.

Suppose f is regular. Let $M(r) := \max_{|z|=r} |f(z)|$.
Then for $r_1 < r < r_2$

$$M(r)^{\log(r_2/r_1)} \leq M(r_1)^{\log(r_2/r)} M(r_2)^{\log(r/r_1)}.$$

Halász Lemma.

For every $k \in \mathbb{N}$, there exists a polynomial
 $h \in \mathcal{P}_k^c$ such that

$$h(0) = 1, \quad h(1) = 0, \quad |h(z)| < \exp\left(\frac{2}{k}\right)$$

for $|z| \leq 1$.

5. PROOFS OF THE MAIN RESULTS

Theorem 2.4. *Every polynomial p of the form*

$$p(x) = \sum_{j=0}^n a_j x^j, \quad |a_n| = 1, \quad |a_j| \leq 1$$

has at most $5\sqrt{n}$ zeros at 1.

Proof of Theorem 2.4. If p has a zero at 1 of multiplicity m , then for every polynomial f of degree less than m , we have

$$(*) \quad a_0 f(0) + a_1 f(1) + \cdots + a_n f(n) = 0.$$

We construct a polynomial f of degree at most $5\sqrt{n}$, for which

$$f(n) > |f(0)| + |f(1)| + \cdots + |f(n-1)|.$$

Equality (*) cannot hold with this f , so the multiplicity of the zero of p at 1 is at most the degree of f .

Let T_ν be the ν -th Chebyshev poly. Let

$$g := T_0 + T_1 + \cdots + T_k \in \mathcal{P}_k.$$

Note that $g(1) = k + 1$ and

$$g(\cos y) = 1 + \cos y + \cos 2y + \cdots + \cos ky$$

$$= \frac{\sin(k + \frac{1}{2})y + \sin \frac{1}{2}y}{2 \sin \frac{1}{2}y}.$$

Hence, for $-1 \leq x < 1$,

$$|g(x)| \leq \frac{\sqrt{2}}{\sqrt{1-x}}.$$

Let $f(x) := g^4\left(\frac{2x}{n} - 1\right)$. Then $f(n) = (k+1)^4$ and

$$|f(0)| + |f(1)| + \cdots + |f(n-1)| \leq \sum_{j=1}^n \frac{4}{\left(\frac{2j}{n}\right)^2} < \frac{\pi^2}{6} n^2.$$

If $k := \lfloor (\pi^2/6)^{1/4} \sqrt{n} \rfloor$ then

$$f(n) > |f(0)| + |f(1)| + \cdots + |f(n-1)|.$$

In this case the degree of f is $4k \leq 5\sqrt{n}$. \square

Theorem 2.5. *For every $n \in \mathbb{N}$, there exists*

$$p_n(x) = \sum_{j=0}^{2n^2} a_j x^j$$

such that $a_{2n^2} = 1$; $a_0, a_1, \dots, a_{2n^2-1}$ are real numbers of modulus less than 1; and p_n has a zero at 1 with multiplicity at least n .

Proof of Theorem 2.5. Define

$$L_n(x) := \frac{(n!)^2}{2\pi i} \int_{\Gamma} \frac{x^t dt}{\prod_{k=0}^n (t - k^2)}$$

where the simple closed contour Γ surrounds the zeros of the denominator in the integrand.

Then L_n is a polynomial of degree n^2 with a zero of order n at 1.

Also, by the residue theorem,

$$L_n(x) = 1 + \sum_{k=1}^n c_{k,n} x^{k^2}$$

where

$$c_{k,n} = \frac{(-1)^n (n!)^2}{\prod_{j=0, j \neq k}^n (k^2 - j^2)} = \frac{(-1)^k 2(n!)^2}{(n-k)!(n+k)!}$$

It follows that

$$c_{k,n} \leq 2, \quad k = 1, 2, \dots, n$$

Hence,

$$q_n(x) := \frac{L_n(x) + L_n(x^2)}{2}$$

is a polynomial of degree $2n^2$ with real coefficients and with a zero at 1 of order n .

Also q_n has constant coefficient 1 and each of its remaining coefficients is a real number of modulus less than 1.

Now let $p_n(x) := x^{2n^2} q_n(1/x)$. \square

Proof of Theorem 2.8. Suppose $P \in \mathcal{A}_n$ has m zeros at -1 . Then $(1+x)^m$ divides P . On evaluating the above at 1 we see that $n \geq 2^m - 1$ and the result follows. \square

6. COMMENTS

- There is an obvious interval dependence in the problem of minimal elements from \mathcal{F}_n .
- On any interval $[0, \delta]$ with $\delta < 1/2$ the only polynomials from \mathcal{F}_n with minimal uniform norm are $\pm x^n$.
- On $[0, 1/2]$ all of $\pm x^n$ and $\pm(x^n - x^{n-1})$ are extremals.
- On any interval $[0, \delta]$ with $\delta > 1/2$ the polynomials $\pm(x^n - x^{n-1})$ work better than x^n , so the nature of the extremals change at $1/2$.

REFERENCES

1. F. Amoroso, *Sur le diamètre transfini entier d'un intervalle réel*, Ann. Inst. Fourier, Grenoble **40** (1990), 885–911.
2. E. Aparicio, *Methods for the approximate calculation of minimum uniform Diophantine deviation from zero on a segment*, Rev. Mat. Hisp.-Amer. **38** (1978), 259–270 (Spanish).
3. E. Aparicio, *New bounds on the minimal Diophantine deviation from zero on $[0, 1]$ and $[0, 1/4]$* , Actus Sextas Jour. Mat. Hisp.-Lusitanas (1979), 289–291.
4. E. Bombieri and J. Vaaler, *Polynomials with low height and prescribed vanishing*, in Analytic Number Theory and Diophantine Problems, Birkhauser (1987), 53–73.
5. P. Borwein and T. Erdélyi, *The integer Cheby-*

- shev problem*, Math. Comp. (to appear).
6. P. Borwein and T. Erdélyi, *Polynomials and Polynomial Inequalities*, Springer-Verlag, New York, 1995.
 7. P. Borwein and C. Ingalls, *The Prouhet, Tarry, Escott problem*, Ens. Math. **40** (1994), 3–27.
 8. Le Baron O. Ferguson, *Approximation by Polynomials with Integral Coefficients*, A.M.S., Rhode Island, 1980.
 9. P. Erdős and P. Turán, *On the distribution of roots of polynomials*, Annals of Math **57** (1950), 105–119.
 10. L.K. Hua, *Introduction to Number Theory*, Springer-Verlag publaddr Berlin Heidelberg, New York, 1982.
 11. J-P Kahane, *Sur les polynômes á coefficients*

- unimodulaires*, Bull. London Math. Soc **12** (1980), 321–342.
12. D.J. Newman and J. S. Byrnes, *The L^4 norm of a polynomial with coefficients ± 1* , MAA Monthly **97** (1990), 42–45.
 13. A. Odlyzko and B. Poonen, *Zeros of polynomials with 0,1 coefficients*, Ens. Math. **39** (1993), 317–348.
 14. G. Pólya and G. Szegő, *Problems and Theorems in Analysis, Volume I*, Springer-Verlag, New York, 1972.
 15. E. B. Saff and R. S. Varga, *On lacunary incomplete polynomials*, Math. Z. **177** (1981), 297–314.
 16. P. Turán, *On a New Method of Analysis and its Applications*, Wiley, New York, 1984.